

CASE REPORTS

13. Cogen FC, Rigg G, Simmons JL, et al: Phencyclidine-associated acute rhabdomyolysis. *Ann Intern Med* 1978 Feb; 88:210-212
14. D'Agostino RS, Arnett EN: Acute myoglobinuria and heroin snorting. *JAMA* 1979 Jan 19; 241:227
15. Browne PM: Rhabdomyolysis and myoglobinuria associated with acute water intoxication. *West J Med* 1979 May; 130:459-461
16. Rheingold OJ, Greenwald RA, Hayes PJ, et al: Myoglobinuria and renal failure associated with typhoid fever. *JAMA* 1977 Jul 27; 238:341
17. Vertel RM, Knochel JP: Acute renal failure due to heat injury: An analysis of ten cases associated with a high incidence of myoglobinuria. *Am J Med* 1967 Sep; 43:435-451
18. Knochel JP, Barcenas C, Cotton JR, et al: Hypophosphatemia and rhabdomyolysis. *Trans Assoc Am Physicians* 1978; 91:156-168
19. Shipley SL, Wrye S: Myoglobinuria. *Heart Lung* 1976 Nov-Dec; 5:950-954
20. Lennette EH: Laboratory diagnosis of viral infections: General principles. *Am J Clin Pathol* 1972 Jun; 57:737-750

Refer to: Chan MKL, Hendrickson CS, Taylor KE: Polyarthrititis associated with breast carcinoma. *West J Med* 1982 Aug; 137:132-133

Polyarthrititis Associated With Breast Carcinoma

MICHAEL KWOK L. CHAN, MD
CHARLES S. HENDRICKSON, MD
KATHLEEN E. TAYLOR, MD
San Francisco

POLYARTHRITIS HAS BEEN REPORTED to be associated with various types of carcinoma.¹⁻⁹ When the primary tumor tissues were removed, the arthritis resolved. Arthritis associated with breast carcinoma was mentioned by MacKenzie and Scherbel,³ who did not, however, describe the nature and extent of the arthritis. Recently, monoarthritis caused by metastatic breast carcinoma was reported by Moutsopoulos and co-workers.⁷ We report the case of a patient with breast carcinoma associated with polyarthrititis and acute inflammation of the synovial fluid. When the primary tumor was removed, the arthritis resolved within a week.

Report of a Case

A 44-year-old East Indian woman without a history of arthritis was admitted to hospital with

a temperature of 37.9°C (100.2°F), malaise and severe incapacitating pain at the right hip and both knees and ankles for three days. A week before the onset of arthritis, she had a needle biopsy of a breast nodule.

At the time she was admitted to hospital, findings on physical examination included a sinus tachycardia of 120 beats per minute with a grade 1 of 6 systolic ejection murmur and a 1.5-cm firm, mobile nodule in the left breast. Tenderness and pain with range of motion were elicited at the right hip. Active synovitis with pain, tenderness, warmth, erythema and moderate swelling was observed at both knees and ankles. Laboratory findings were as follows: a leukocyte count of 11,300 per cu mm with no left shift, a hematocrit of 38 percent and an erythrocyte sedimentation rate (Westergren method) of 90 mm per hr (repeat testing showed 148 mm per hr). The results of electrolyte, liver function, rheumatoid factor and antinuclear antibody (ANA) tests were all normal. Analysis of knee aspirate gave the following values: leukocyte count, 65,000 per cu mm, with 94 percent polymorphonuclear leukocytes; glucose, 111 mg per dl; protein, 5 grams per dl; C₃, 81 mg per dl; no crystals; negative findings on acid-fast smear, and negative cultures for *Neisseria gonorrhoeae* and *Brucella* sp. Cultures of specimens from throat and cervix showed normal flora; the results of blood cultures were negative. The patient was given penicillin and nafcillin intravenously but continued to have temperatures up to 39.6°C (103.3°F) with nausea and vomiting. No apparent improvement occurred in her joint symptoms, and antibiotic therapy was discontinued. The fever abated, however, and she was discharged on a regimen of indomethacin.

The joint symptoms lessened slightly during the next few weeks, then recurred. Active synovitis with symmetric distribution now affected the proximal interphalangeal joints, metacarpophalangeal joints, wrists, elbows, knees and ankles. X-ray films of the sacroiliac joint and hips were normal. She was treated with naproxen and given an intramuscular injection of dexamethasone; her condition improved moderately. The biopsy specimen of the breast nodule showed medullary carcinoma, for which mastectomy was carried out. No lymph node involvement was found. Naproxen therapy was discontinued postoperatively. Three days after the operation, her joint symptoms de-

From the Departments of Medicine, Kaiser-Permanente Medical Center, San Francisco, and Stanford University Medical Center, Stanford, California. Dr. Chan is now in private practice.

Submitted, revised, July 30, 1981.

Reprint requests to: Michael Kwok L. Chan, MD, Department of Internal Medicine-Rheumatology, Oak Hill Medical Group, 3100 Telegraph Avenue, Oakland, CA 94609.

creased considerably; by the seventh postoperative day, all joint symptoms had resolved. At multiple follow-up visits over the next 12 months, the patient showed no evidence of arthritis.

Discussion

Various musculoskeletal symptoms have been described as the presenting features of an underlying malignant condition. These symptoms, ranging from myalgia and arthralgia to clinically evident monoarthritis or polyarthritis, all resolve when the primary tumor is removed. In our patient, a rapid and disabling polyarthritis developed, which was only partially controlled with nonsteroidal antiinflammatory drugs. When a breast tumor was removed, the arthritis resolved within a week.

The data in the literature on synovial fluid analysis in patients with a malignant lesion have been sparse. Noninflammatory fluid has been described by Ropes and Bauer¹⁰ in patients with pulmonary hypertrophic osteoarthropathy and in the patients reported by Moutsopoulos and associates.⁷ In our patient, analysis of a specimen of the synovial fluid from the knee showed an acute inflammatory reaction. Zacharski and Kyle¹¹ reported that the erythrocyte sedimentation rate was greatly elevated (more than 100 mm an hour, Westergren method) in patients with malignancy. In our patient, the erythrocyte sedimentation rate was elevated to 148 mm per hour (Westergren method).

The pathogenesis of arthritis in association with malignancy is unknown. Husain and colleagues¹² showed a positive result on ANA test to be associated with carcinoma. Sheon and associates¹³ reported a positive rheumatoid factor in 27 percent of patients with malignancy. In Bartfeld's¹⁴ survey the incidence of positive rheumatoid factor in patients with malignancy ranged from 1 percent to 20 percent. Circulating immune complexes and hypocomplementemia in patients with carcinoma were reported by Teshima and co-workers¹⁵ and by Rosen and colleagues.¹⁶ Recently, the prostaglandin E₂ (PGE₂) level was shown to increase in patients with metastatic bone disease.^{17,18} Pretreatment with administration of indomethacin reduces the amount of PGE₂ extractable from the tumor, as has been proved by Galasko¹⁹ and by Voelkel and associates.²⁰ Bennett and co-workers²¹ and Easty and Easty²² separately found an increased level of PGE₂ in breast carcinoma tis-

sues. This finding was recently confirmed by Roland and colleagues.²³ Prostaglandin E₂ is known to be one of the mediators of inflammation. Whether this local increase in PGE₂ relates causally or temporally to the development of arthritis needs further investigation.

Summary

Polyarthritis associated with several types of carcinoma has been reported. We report the case of a patient with breast carcinoma in whom polyarthritis resolved when the tumor was removed. An increase of the amount of prostaglandin E₂ in breast carcinoma tissue, which has been reported in the literature, might be related to the development of arthritis.

REFERENCES

1. Lansbury J: Collagen disease complicating malignancy. *Ann Rheum Dis* 1953; 12:301-305
2. Harrison JW, Scherbel AL: Rheumatic musculoskeletal symptoms associated with carcinoma of the prostate gland. *Postgrad Med* 1960; 28:274-283
3. MacKenzie AH, Scherbel AL: Connective tissue syndromes associated with carcinoma. *Geriatrics* 1963; 18:745-753
4. Calabro JJ: Cancer and arthritis. *Arthritis Rheum* 1967; 10:553-567
5. Virshup AM, Sliwiski AJ: Polyarthritis and subcutaneous nodules associated with carcinoma of the pancreas. *Arthritis Rheum* 1973; 16:388-391
6. Goldenberg DL, Kelley W, Gibbons RB: Metastatic adenocarcinoma of synovium presenting as an acute arthritis—Diagnosis by closed synovial biopsy. *Arthritis Rheum* 1975; 18:107-110
7. Moutsopoulos HM, Fye KH, Pugay PI, et al: Monoarthritic arthritis caused by metastatic breast carcinoma—Value of cytologic study of synovial fluid. *JAMA* 1975 Oct 6; 234:75-76
8. Meals RA, Hungerford DS, Stevens MB: Malignant disease mimicking arthritis of the hip. *JAMA* 1978 Mar 13; 239:1070-1071
9. Simon RD Jr, Ford LE: Rheumatoid-like arthritis associated with a colonic carcinoma. *Arch Intern Med* 1980; 140:698-700
10. Ropes MW, Bauer W: *Synovial Fluid Changes in Joint Disease*. Cambridge, Harvard University Press, 1953
11. Zacharski LR, Kyle RA: Significance of extreme elevation of erythrocyte sedimentation rate. *JAMA* 1967; 202:264-266
12. Husain M, Neff J, Daily E, et al: Antinuclear antibodies: Clinical significance of titers and fluorescence patterns. *Am J Clin Pathol* 1974; 61:59-65
13. Sheon RP, Kirsner AB, Tangintanapap P, et al: Malignancy in rheumatic disease: Interrelationships. *J Am Geriatr Soc* 1977; 25:20-27
14. Bartfeld H: Incidence and significance of seropositive tests for rheumatoid factor in non-rheumatoid diseases. *Ann Intern Med* 1960; 52:1059-1066
15. Teshima H, Wanebo H, Pinsky C, et al: Circulating immune complexes detected by ¹²⁵I-Clq deviation test in sera of cancer patients. *J Clin Invest* 1977; 59:1134-1142
16. Rosen RD, Barnes BC: Measuring serum immune complexes in cancer. *Ann Intern Med* 1978; 88:570-571
17. Bennett A, McDonald AM, Simpson JS, et al: Breast cancer, prostaglandins, and bone metastases. *Lancet* 1975; 1:1218-1220
18. Bennett A, Charlier EM, McDonald AM, et al: Bone destruction by breast tumors. *Prostaglandins* 1976; 11:461-463
19. Galasko CSB: Relationship of bone destruction in skeletal metastases to osteoclast activation and prostaglandins. *Nature* 1976; 263:508-510
20. Voelkel EF, Tashjian AH Jr, Franklin R, et al: Hypercalcemia and tumor-prostaglandins: The VX₂ carcinoma model in the rabbit. *Metabolism* 1975; 24:973-986
21. Bennett A, Charlier EM, McDonald AM, et al: Prostaglandins and breast cancer. *Lancet* 1977; 2:624-626
22. Easty GC, Easty DM: Prostaglandins and cancer. *Cancer Treat Rev* 1976; 3:217-225
23. Rolland PH, Martin PM, Jacquemier J, et al: Prostaglandin production and metabolism in human breast cancer. In Samuelsen B, Ramwell PW, Paoletti R (Eds): *Advances in Prostaglandin and Thromboxane Research*, Vol 6. New York, Raven, 1980, pp 574-580